

KEK-TH-329
 KEK Preprint 92-27
 April 1992

Fusion Rules for the q -Vertex Operators of $U_q(\widehat{sl(2)})$

HIDETOSHI AWATA ¹ and YASUHIKO YAMADA ²

^{1, 2} *National Laboratory for High Energy Physics (KEK)*

Tsukuba, Ibaraki 305, Japan

and

¹*Dept. of Physics Hokkaido University,*

Sapporo 060, Japan

Abstract

We analyze the properties of the q -vertex operators of $U_q(\widehat{sl(2)})$ introduced by Frenkel and Reshetikhin. As the condition for the null vector decoupling, we derive the existence condition of the q -vertex operators (the fusion rules).

¹ E-mail address : awata@jpnkekvm.bitnet

² E-mail address : yamaday@jpnkekvm.bitnet

1. Introduction

The q -deformed chiral vertex operators for the quantum affine Lie algebra $U_q(\widehat{\mathfrak{g}})$, defined by Frenkel and Reshetikhin [1], are the q -deformed version of the ordinary chiral vertex operators of the WZNW model introduced by Tsuchiya and Kanie [2], and they have remarkable applications to the solvable lattice models [3, 4]. In the present paper, we investigate the fusion rules for the q -vertex operators of $U_q(\widehat{sl(2)})$.

This paper is arranged as follows. In section 2, we briefly summarize the $U_q(\widehat{sl(2)})$ algebra. In section 3, we introduce the q -vertex operators. We discuss their matrix elements and q -operator product expansion. Next, in section 4, we present the null vectors explicitly and calculate the matrix elements of the q -vertex operator including the null vectors. We give the fusion rules in section 5. This paper is a q -analogue of our previous one [5].

2. Quantum Algebra $U_q(\widehat{sl(2)})$

§ **2.1.** First we define some notation. The algebra $U_q(\widehat{sl(2)})$ is generated by e_i , f_i and invertible k_i ($i = 0, 1$) with relations

$$\begin{aligned}
 k_i e_j k_i^{-1} &= q^{a_{ij}} e_j, & k_i f_j k_i^{-1} &= q^{-a_{ij}} f_j, \\
 e_i f_j - f_j e_i &= \delta_{ij} \frac{k_i - k_i^{-1}}{q - q^{-1}}, \\
 \sum_{n=0}^{1-a_{ij}} (-1)^n \begin{bmatrix} 1-a_{ij} \\ n \end{bmatrix} e_i^{1-a_{ij}-n} e_j e_i^n &= 0 & (i \neq j) \\
 \sum_{n=0}^{1-a_{ij}} (-1)^n \begin{bmatrix} 1-a_{ij} \\ n \end{bmatrix} f_i^{1-a_{ij}-n} f_j f_i^n &= 0 & (i \neq j)
 \end{aligned} \tag{2.1}$$

where

$$\begin{aligned} a_{ij} &= \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}, & \begin{bmatrix} n \\ m \end{bmatrix} &= \frac{[n]!}{[n-m]![m]!}, \\ [n]! &= [n] \cdots [1], & [n] &= \frac{q^n - q^{-n}}{q - q^{-1}}. \end{aligned} \quad (2.2)$$

The algebra $U_q(\widehat{sl(2)})$ is a Hopf algebra with the comultiplication Δ , the antipode S and the co-unit ϵ

$$\begin{aligned} \Delta k_i &= k_i \otimes k_i, \\ \Delta e_i &= e_i \otimes k_i + 1 \otimes e_i, & \Delta f_i &= f_i \otimes 1 + k_i^{-1} \otimes f_i, \\ S(e_i) &= -e_i k_i^{-1}, & S(k_i) &= k_i^{-1}, & S(f_i) &= -k_i f_i, \\ \epsilon(e_i) &= \epsilon(f_i) = 0, & \epsilon(k_i) &= 1. \end{aligned} \quad (2.3)$$

There exists a difference analog of the Virasoro L_0 operator $l_0 (= q^{2(k+2)L_0} = p^{L_0}) \in U_q(\widehat{sl(2)})$ such that $l_0 k_i = k_i l_0$, $l_0 e_i = p^{-\delta_{i,0}} e_i l_0$ and $l_0 f_i = p^{\delta_{i,0}} f_i l_0$, where $p = q^{2(k+2)}$.

§ 2.2. Let M_j be the Verma module over $U_q(\widehat{sl(2)})$, generated by the highest weight vector $|j\rangle$, such that $e_i |j\rangle = 0$, $k_1 |j\rangle = q^{2j} |j\rangle$ and $k_0 |j\rangle = q^{k-2j} |j\rangle$. Here $k \in \mathbf{C}$ is a level. Instead of k , sometimes we use $t (= k + 2)$. The action of l_0 on M_j is defined by $l_0 |j\rangle = p^{h_j} |j\rangle$ with $h_j = j(j+1)/(k+2)$.

The dual module M_j^* is generated by $\langle j|$ which satisfies $\langle j|e_i = 0$, $\langle j|k_1 = q^{2j} \langle j|$ and $\langle j|k_0 = q^{k-2j} \langle j|$. The bilinear form $M_j^* \otimes M_j \rightarrow \mathbf{C}$ is uniquely defined by $\langle j|j\rangle = 1$ and $(\langle u|X)|v\rangle = \langle u|(X|v\rangle)$ for any $\langle u| \in M_j^*$, $|v\rangle \in M_j$ and $X \in U_q(\widehat{sl(2)})$.

For a generic value of q , the representation theory of $U_q(\widehat{sl(2)})$ is the same as that of $\widehat{sl(2)}$, and there are also the null vectors, which control the reducibility of the representation. A null vector $|\chi\rangle \in M_j$ (of grade N and charge Q) is defined by $e_i |\chi\rangle = 0$, $k_1 |\chi\rangle = q^{2(j+Q)} |\chi\rangle$, $k_0 |\chi\rangle = q^{k-2(j+Q)} |\chi\rangle$ and $l_0 |\chi\rangle = p^{h_j+N} |\chi\rangle$. A null vector $\langle \chi| \in M_j^*$ is defined in a similar manner.

Example. Let us give some examples of the null vectors. Set $2j_{r,s} + 1 = r - st$, $r, s \in \mathbf{Z}$ and $t = k + 2 \in \mathbf{C}$. Examples of the null vectors $|\chi_{r,s}\rangle$ are,

$$\begin{aligned} |\chi_{r,0}\rangle &= (f_1)^r |j_{r,0}\rangle \quad \text{for } r > 0 \\ |\chi_{r,0}\rangle &= (f_0)^{-r} |j_{r,0}\rangle \quad \text{for } r < 0 \\ |\chi_{1,1}\rangle &= \left(\frac{f_0 f_1 f_1}{[t+1]} - \frac{[2] f_1 f_0 f_1}{[t]} + \frac{f_1 f_1 f_0}{[t-1]} \right) |j_{1,1}\rangle. \end{aligned} \quad (2.4)$$

It is easy to show $e_1|\chi\rangle = 0$ and $e_0|\chi\rangle = 0$. For example,

$$e_0|\chi_{1,1}\rangle = \left(\frac{[2t+2]}{[t+1]} - \frac{[2][2t]}{[t]} + \frac{[2t-2]}{[t-1]} \right) f_1 f_1 |j_{1,1}\rangle = 0, \quad (2.5)$$

owing to the following relation ; $\frac{[a]}{[b]} - \frac{[2][a+2]}{[b+1]} + \frac{[a+4]}{[b+2]} = \frac{[2][a-2b]}{[b][b+1][b+2]}.$

3. The q -deformed Vertex Operator

§ 3.1. For a general Hopf algebra U , the field operator $\phi(z)$ is defined by the following transformation property under the adjoint action of U : [1, 6, 7]

$$\text{ad}(a)\phi(z) \equiv \sum_k a_k^{(1)} \phi(z) S(a_k^{(2)}) = \rho(a)\phi(z), \quad (3.1)$$

where $a \in U$, $\Delta^{(n)}(a) = \sum_k a_k^{(1)} \otimes \cdots \otimes a_k^{(n)}$ is the comultiplication $\Delta^{(n)} : U \rightarrow U^{\otimes n}$, and ρ is a certain representation of U . The adjoint action is the natural generalization of the commutator for Lie algebras.

For $U = U_q(\widehat{sl(2)})$, the q -vertex operator $\phi_j(z, x)$ of spin j is defined explicitly as follows

$$\begin{aligned} k_i \phi_j(z, x) &= \rho(k_i) \phi_j(z, x) k_i, \\ e_i \phi_j(z, x) &= \rho(e_i) \phi_j(z, x) k_i + \phi_j(z, x) e_i, \\ f_i \phi_j(z, x) &= \rho(f_i) \phi_j(z, x) + \rho(k_i^{-1}) \phi_j(z, x) f_i, \end{aligned}$$

and

$$l_0 \phi_j(z, x) = \rho(l_0) \phi_j(z, x) l_0, \quad (3.2)$$

where $\rho = \rho_{j,x}$ is a contravariant representation defined by

$$\begin{aligned}\rho(k_1) &= q^{2(j-x\frac{d}{dx})}, & \rho(k_0) &= q^{-2(j-x\frac{d}{dx})}, \\ \rho(e_1) &= x[2j - x\frac{d}{dx}], & \rho(e_0) &= \frac{z}{x}[x\frac{d}{dx}], \\ \rho(f_1) &= \frac{1}{x}[x\frac{d}{dx}], & \rho(f_0) &= \frac{x}{z}[2j - x\frac{d}{dx}],\end{aligned}\tag{3.3}$$

$$\rho(l_0) = p^{z\frac{d}{dz}}.$$

Our definition of the q -vertex operator is useful in actual calculations.

In the case of $2j \in \mathbf{Z}_{\geq 0}$, $\rho(k_1)$, $\rho(e_1)$ and $\rho(f_1)$ give the $2j + 1$ dimensional representation of $U_q(sl(2))$ on $\oplus_{m=0}^{2j} \mathbf{C}x^m$.

§ 3.2. We will investigate the q -vertex operator $\phi_{j_2}(z, x)$ in the following two interpretations: (1) the matrix element and (2) the operator.

(1) Matrix element $\phi : M_{j_3}^* \otimes M_{j_1} \rightarrow \mathbf{C}$.

From the k_1 and l_0 commutation relations, the ground state matrix element of the q -vertex operator is given by

$$\langle j_3 | \phi_{j_2}(z, x) | j_1 \rangle = C_{123} z^{h_{13}} x^{j_{13}},\tag{3.4}$$

where $h_{13} = -h_1 + h_3$, $j_{13} = j_1 + j_2 - j_3$ and C_{123} is an arbitrary constant.

The other matrix elements for the descendant fields are uniquely determined by the $U_q(\widehat{sl(2)})$ algebra. For example,

$$\begin{aligned}\langle j_3 | \phi_{j_2}(z, x) f_{\alpha_1} \cdots f_{\alpha_n} | j_1 \rangle &= (-1)^n \rho(k_{\alpha_1}) \rho(f_{\alpha_1}) \cdots \rho(k_{\alpha_n}) \rho(f_{\alpha_n}) \langle j_3 | \phi_{j_2}(z, x) | j_1 \rangle, \\ \langle j_3 | e_{\alpha_1} \cdots e_{\alpha_n} \phi_{j_2}(z, x) | j_1 \rangle &= \rho(e_{\alpha_1}) \cdots \rho(e_{\alpha_n}) \langle j_3 | \phi_{j_2}(z, x) k_{\alpha_1} \cdots k_{\alpha_n} | j_1 \rangle.\end{aligned}\tag{3.5}$$

Note that, if $C_{123} = 0$, then $\phi_{j_2}(z, x) = 0$.

(2) Operator $\phi : M_{j_1} \rightarrow M_{j_3}$.

To study the property of the map $\phi : M_{j_1} \rightarrow M_{j_3}$, let us consider the image of the ground state $|j_1\rangle \in M_{j_1}$ (the q -operator product expansion),

$$\phi_{j_2}(z, x)|j_1\rangle = \sum_{N, Q} \sum_J \beta_J^{NQ} q^{kN-2j_1Q} z^{h_{13}+N} x^{j_{13}-Q} |J^{NQ}\rangle_{j_3}, \quad (3.6)$$

where $|J^{NQ}\rangle_{j_3} \in M_{j_3}$ is the basis of the homogeneous components of grade N and charge Q .

The coefficients $\beta_J = \beta_J^{NQ}$, for fixed N, Q in (3.6), are determined by the following linear equations,

$$\sum_J \langle I|J\rangle \beta_J = \langle I|\phi_{j_2}(z, x)|j_1\rangle \Big|_{N, Q}, \quad (3.7)$$

where $\Big|_{N, Q}$ denotes the coefficient of $q^{kN-2j_1Q} z^{h_{13}+N} x^{j_{13}-Q}$. In the Appendix, we give some sample calculations of the coefficients β .

The vectors $|N, Q\rangle = \sum_J \beta_J^{NQ} |J^{NQ}\rangle_{j_3}$ are also characterized by the descent equations

$$\begin{aligned} e_1|N, Q\rangle &= [-j_1 + j_2 + j_3 + Q + 1]|N, Q + 1\rangle, \\ e_0|N, Q\rangle &= [j_1 + j_2 - j_3 - Q + 1]|N - 1, Q - 1\rangle. \end{aligned} \quad (3.8)$$

This is the q -analogue of the descent equations in [8].

§ 3.3. If the determinant of the matrix $\langle I|J\rangle$ (the Shapovalov form) vanishes, the consistency of the equation (3.7) requires the following constraints,

$$\langle \chi|\phi_{j_2}(z, x)|j_1\rangle = 0, \quad (3.9)$$

for any null vector $\langle \chi| \in M_{j_3}^*$. This is the so-called the null vector decoupling condition. Similarly, consistency of the q -OPE of $\phi : M_{j_3}^* \rightarrow M_{j_1}^*$, i.e. $\langle j_3|\phi_{j_2}(z, x)$, requires $\langle j_3|\phi_{j_2}(z, x)|\chi\rangle = 0$ for all null vectors $|\chi\rangle \in M_{j_1}$. Therefore, the existence of the q -vertex operator depends on the null vector decoupling.

Example. Here we give some examples of the null vector decoupling condition.

(i). For $2j_1 + 1 = r \in \mathbf{Z}_{>0}$, by using the null vector $|\chi_{r,0}\rangle$ in (2.4),

$$\langle j_3 | \phi_{j_2} | \chi_{r,0} \rangle = (-1)^r \frac{[j_{13}]!}{[j_{13} - r]!} q^{(2j_3 - 2j_1 + r + 1)r} z^{h_{13} - r} x^{j_{13}}. \quad (3.10)$$

Hence, the null vector decoupling condition is $\prod_{n=0}^{r-1} [j_1 + j_2 - j_3 - n] = 0$.

(ii). For $2j_1 + 1 = 1 - t$,

$$\begin{aligned} \langle j_3 | \phi_{j_2} | \chi_{1,1} \rangle &= - \left(\frac{[2j_2 - j_{13} + 2][j_{13} - 1][j_{13}]}{[t + 1]} - \frac{[2][j_{13}][2j_2 - j_{13} + 1][j_{13}]}{[t]} \right. \\ &\quad \left. + \frac{[j_{13}][j_{13} + 1][2j_2 - j_{13}]}{[t - 1]} \right) q^{2j_3 - 2j_1 + 4} z^{h_{13} - 1} x^{j_{13} - 1} \\ &= - \frac{[2][2j_2 - j_{13} + t][j_{13} + 1 - t][2j_2 - j_{13}]}{[t + 1][t][t - 1]} q^{2j_3 - 2j_1 + 4} z^{h_{13} - 1} x^{j_{13} - 1}. \end{aligned} \quad (3.11)$$

Here we use the following formula ; $\frac{[a][b]}{[c]} - \frac{[2][a+1][b+1]}{[c+1]} + \frac{[a+2][b+2]}{[c+2]} = \frac{[2][a-c][b-c]}{[c][c+1][c+2]}$.

Therefore, the null vector decoupling condition is

$$[j_3 + j_2 - j_1 + t][j_1 + j_2 - j_3 + 1 - t][j_3 + j_2 - j_1] = 0. \quad (3.12)$$

4. The Null Vectors

Here we analyze the null vector decoupling condition by using the general form of the null vectors which is given by Malikov [9].

For $t = k + 2 \in \mathbf{C} \setminus \{0\}$ and the highest weight j , parametrized as $2j_{r,s} + 1 = r - st$ with $r, s \in \mathbf{Z}$, such that (i) $r > 0$ and $s \geq 0$ or (ii) $r < 0$ and $s < 0$, there exists a unique null vector $|\chi_{r,s}\rangle \in M_j$ of grade $N = rs$ and charge $Q = -r$. And the null vector in $M_{j_{r,s}}$ is as follows,

$$\begin{aligned} |\chi_{r,s}\rangle &= (f_1)^{r+st} (f_0)^{r+(s-1)t} \dots (f_0)^{r-(s-1)t} (f_1)^{r-st} |j_{r,s}\rangle, \\ |\chi_{r,s}\rangle &= (f_0)^{-r-(s+1)t} (f_1)^{-r-(s+2)t} \dots (f_1)^{-r+(s+2)t} (f_0)^{-r+(s+1)t} |j_{r,s}\rangle, \end{aligned} \quad (4.1)$$

for the cases (i) and (ii) respectively. These formulae make sense after analytic continuation.

We now consider the matrix element including the null vector $|\chi_{r,s}\rangle \in M_{j_1}$. From (3.2), (3.4) and (4.1), we obtain

$$\langle j_3 | \phi_{j_2}(z, x) | \chi_{r,s} \rangle = (-1)^{(2s+1)r} C_{123} f_{r,s}(j_1, j_2, j_3) q^{(2j_3-2j_1+r+2s+1)r} z^{h_{13}-rs} x^{j_{13}-r}, \quad (4.2)$$

where

$$\begin{aligned} f_{r,s}(j_1, j_2, j_3) &= \prod_{n=0}^{r-1} \prod_{m=0}^s [j_1 + j_2 - j_3 - n + mt] \prod_{n=1}^r \prod_{m=1}^s [-j_1 + j_2 + j_3 + n - mt], \\ f_{r,s}(j_1, j_2, j_3) &= \prod_{n=0}^{-r-1} \prod_{m=0}^{-s-1} [-j_1 + j_2 + j_3 - n + mt] \prod_{n=1}^{-r} \prod_{m=1}^{-s-1} [j_1 + j_2 - j_3 + n - mt]. \end{aligned} \quad (4.3)$$

for the cases (i) and (ii) respectively. The proof is given by a method similar to that used in [5]. Here we use, for instance,

$$\left(\frac{1}{x}[x\partial]\right)^n x^j = \frac{[j]!}{[j-n]!} x^{j-n},$$

for any $n, j \in \mathbf{C}$. The formulae (4.3) are our main results. Note that similar expressions are obtained for a projection of the singular vectors [9].

Next, we consider another matrix element $\langle \chi_{r,s} | \phi_{j_2}(z, x) | j_1 \rangle$. This element follows from $\langle j_3 | \phi_{j_2}(z, x) | \chi_{r,s} \rangle$, because of the existence of the following anti-algebra automorphism. Indeed, the σ defined by

$$\begin{aligned} \sigma(e_i) &= -q^{-2} f_i k_i, \quad \sigma(k_i) = k_i, \quad \sigma(f_i) = -q^2 k_i^{-1} e_i, \\ \sigma(dx^{-j} \phi_j(z, x)) &= dy^{-j} \phi_j(w, y), \quad \sigma(dx^{-j} \langle u | \phi_j(z, x) | v \rangle) = dx^{-j} \langle u | \phi_j(z, x) | v \rangle, \end{aligned} \quad (4.4)$$

with $w = z^{-1}$ and $y = x^{-1}$, is the anti-algebra automorphism, which means $\sigma(k_i^{-1}) \sigma(e_j) \sigma(k_i) = q^{a_{ij}} \sigma(e_j)$, $\Delta \sigma(e_i) = \sigma(e_i) \otimes \sigma(k_i) + \sigma(1) \otimes \sigma(e_i)$ and $\sigma(\phi_j(z, x)) \sigma(k_i) = \sigma(k_i) \rho(k_i) \sigma(\phi_j(z, x))$ etc. Consequently, we obtain

$$\langle \chi_{r,s} | \phi_{j_2}(z, x) | j_1 \rangle = (-1)^{(2s+1)r} C_{123} f_{r,s}(j_3, j_2, j_1) q^{(2j_1-2j_3+r+2s+1)r} z^{h_{13}+rs} x^{j_{13}+r}, \quad (4.5)$$

with the same function $f_{r,s}(j_3, j_2, j_1)$ as (4.3).

5. Fusion Rules

The factor $f_{rs}(j_1, j_2, j_3)$ is written in terms of the q -integer; however, this factor is almost the same as that of the $\widehat{sl}(2)$ case [5]. Therefore, the fusion rules are similar to the $\widehat{sl}(2)$ case, if q is not a root of unity.

§ 5.1. When M_{j_1} has a null vector $|\chi_{r,s}\rangle$, there exists a q -vertex operator if and only if the matrix element including the null vector vanishes, $\langle j_3 | \phi_{j_2}(z, x) | \chi_{r,s} \rangle = 0$, i.e. $f_{r,s}(j_1, j_2, j_3) = 0$. And this gives the fusion rules.

In the case that $M_{j_3}^*$ also has a null vector $\langle \chi_{r_3, s_3} |$, the existence condition of the q -vertex operator is $\langle j_3 | \phi_{j_2}(z, x) | \chi_{r_1, s_1} \rangle = \langle \chi_{r_3, s_3} | \phi_{j_2}(z, x) | j_1 \rangle = 0$, i.e. $f_{r_1, s_1}(j_1, j_2, j_3) = f_{r_3, s_3}(j_3, j_2, j_1) = 0$.

§ 5.2. If the level is rational $t = p/q$, with the coprime integers p and q , then $j_{r,s} = j_{r-p, s-q}$. Hence, there are two independent null vectors for $j_1 = j_{r,s} = j_{r-p, s-q}$. There exists a q -vertex operator if and only if $\langle j_3 | \phi_{j_2}(z, x) | \chi_{r,s} \rangle = \langle j_3 | \phi_{j_2}(z, x) | \chi_{r-p, s-q} \rangle = 0$, i.e. $f_{r,s}(j_1, j_2, j_3) = f_{r-p, s-q}(j_1, j_2, j_3) = 0$.

The last fusion rule says that M_{j_3} also has two null vectors. Hence, we also need the condition $\langle \chi_{r,s} | \phi_{j_2}(z, x) | j_1 \rangle = \langle \chi_{r-p, s-q} | \phi_{j_2}(z, x) | j_1 \rangle = 0$, that is $f_{r,s}(j_3, j_2, j_1) = f_{r-p, s-q}(j_3, j_2, j_1) = 0$. However, this condition is included in the above one.

In Ref [3], it is shown that the space of vertex operators has a basis indexed by admissible triples $\{\Phi_\lambda^{\mu b}(z)\}$, for dominant integral weights λ and μ for any $U_q(\widehat{\mathfrak{g}})$. For the integrable case, i.e. k is a non-negative integer, our results agree with those of Ref [3].

6. Conclusions and Remarks

As the condition for the null vector decoupling, we have derived the fusion rules for the $U_q(\widehat{sl(2)})$ algebra. Similar results are expected for other quantum affine Lie algebras. When q is a root of unity, the structure of the representations changes drastically, and we need further investigation to understand the q -vertex operators in such cases.

Acknowledgments

We would like to thank F. G. Malikov, A. Tsuchiya and the members of KEK theory group for valuable discussions. We would also like to thank to N. A. McDougall for a careful reading of the manuscript.

Appendix

Let us give some examples of the q -OPE calculation. Set $a \equiv k - 2j_3$, $b \equiv 2j_3$ and $j_{13} \equiv j_1 + j_2 - j_3$. In (3.6), let

$$\sum_J \beta_J^{NQ} |J^{NQ}\rangle_{j_3} = \sum_{\alpha_1, \dots, \alpha_n} \beta_{\alpha_1, \dots, \alpha_n} f_{\alpha_1} \cdots f_{\alpha_n} |j_3\rangle \quad (\text{A.1})$$

where $N = \sum_i \bar{\alpha}_i$, $Q = \sum_i \bar{\alpha}_i - \sum_i \alpha_i$, $\alpha_i = 0, 1$ and $\bar{\alpha} = 1 - \alpha$.[†]

(i). For $N = 0$ and $Q = -r \in \mathbf{Z}$, the basis is $(f_1)^r |j_3\rangle$, the Shapovalov form is $\langle j_3 | (e_1)^r (f_1)^r |j_3\rangle = [r]! [b]! / [b - r]!$ and $\langle j_3 | (e_1)^r \phi_{j_2}(z, x) |j_1\rangle = q^{2j_1 r} z^{h_{13}} x^{j_{13} + r} [2j_2 - j_{13}]! / [2j_2 - j_{13} - r]!$. Hence, we obtain

$$\frac{[r]! [b]!}{[b - r]!} \underbrace{\beta_1, \dots, 1}_{r \text{ times}} = \frac{[2j_2 - j_{13}]!}{[2j_2 - j_{13} - r]!}. \quad (\text{A.2})$$

If $[b - r + 1] = [2j_3 - r + 1] = 0$, then $\langle j_3 | (e_1)^r$ is a null vector; therefore the existence condition of the q -vertex operator $\phi_{j_2} : M_{j_1} \rightarrow M_{j_3}$ is $\prod_{n=0}^{r-1} [j_3 + j_2 - j_1 - n] = 0$.

[†] In general, such a basis is an over complete one.

(ii). For $N = 1$ and $Q = 0$, the basis is $f_1 f_0 |j_3\rangle$ and $f_0 f_1 |j_3\rangle$ and the Shapovalov form is

$$\begin{pmatrix} \langle j_3 | e_1 e_0 f_1 f_0 | j_3 \rangle & \langle j_3 | e_1 e_0 f_0 f_1 | j_3 \rangle \\ \langle j_3 | e_0 e_1 f_1 f_0 | j_3 \rangle & \langle j_3 | e_0 e_1 f_0 f_1 | j_3 \rangle \end{pmatrix} = \begin{pmatrix} [a][b] & [a+2][b] \\ [a][b+2] & [a][b] \end{pmatrix}. \quad (\text{A.3})$$

From (3.7), we can obtain the β

$$[2][a][b][a+b+2] \begin{pmatrix} \beta_{10} \\ \beta_{01} \end{pmatrix} = \begin{pmatrix} -[a][b] & [a+2][b] \\ [a][b+2] & -[a][b] \end{pmatrix} \begin{pmatrix} [j_{13}+1][2j_2-j_{13}] \\ [2j_2-j_{13}+1][j_{13}] \end{pmatrix}. \quad (\text{A.4})$$

(iii). For $N = 1$ and $Q = -1$, the basis is $f_1 f_1 f_0 |j_3\rangle$, $f_1 f_0 f_1 |j_3\rangle$ and $f_0 f_1 f_1 |j_3\rangle$, and the Shapovalov form is

$$[2] \begin{pmatrix} [a][b-1][b] & [a+2][b-1][b] & [a+4][b-1][b] \\ [a][b][b+1] & [a+1][b][b] & [a+2][b-1][b] \\ [a][b+1][b+2] & [a][b][b+1] & [a][b-1][b] \end{pmatrix}. \quad (\text{A.5})$$

Therefore, we have

$$[2]^2 [a][b][b-1][a+b+2][a+2b+2] \begin{pmatrix} \beta_{110} \\ \beta_{101} \\ \beta_{011} \end{pmatrix} = \begin{pmatrix} [a][b][b-1] & -[2][a][b][b-1] & [b][b-1] \\ \times [a+b+2] & \times [a+b+3] & \times \begin{pmatrix} [2][a+2][a+b+3] \\ -[a+4][a+b+2] \end{pmatrix} \\ -[2][a][b-1][b+1] & [2][a][b-1] & -[2][a][b][b-1] \\ \times [a+b+2] & \times \begin{pmatrix} [b-1][a+b+2] \\ +[b+2][a+b+3] \end{pmatrix} & \times [a+b+3] \\ [a][b][b+1] & -[2][a][b-1][b+1] & [a][b][b-1] \\ \times [a+b+2] & \times [a+b+2] & \times [a+b+2] \end{pmatrix}$$

$$\times \begin{pmatrix} [j_{13} + 2][2j_2 - j_{13} - 1][2j_2 - j_{13}] \\ [2j_2 - j_{13}][j_{13} + 1][2j_2 - j_{13}] \\ [2j_2 - j_{13}][2j_2 - j_{13} + 1][j_{13}] \end{pmatrix}. \quad (\text{A.6})$$

If the determinant of the Shapovalov form vanishes, i.e. $[a + 2b + 2] = [k + 2j_3 + 2] = 0$, then there is a null vector $\langle \chi |$ such as

$$\langle \chi | = \langle j_3 | \left(\frac{e_1 e_1 e_0}{[t + 1]} - \frac{[2] e_1 e_0 e_1}{[t]} + \frac{e_0 e_1 e_1}{[t - 1]} \right). \quad (\text{A.7})$$

Hence the existence condition of the q -vertex operator is $[j_1 + j_2 - j_3 + t][j_3 + j_2 - j_1 + 1 - t][j_1 + j_2 - j_3] = 0$.

(iv). Finally, there is a following relation

$$\beta_{\alpha_1, \dots, \alpha_n}(j_1, j_2, j_3) = \beta_{\bar{\alpha}_1, \dots, \bar{\alpha}_n}\left(\frac{k}{2} - j_1, j_2, \frac{k}{2} - j_3\right). \quad (\text{A.8})$$

References

1. I.B. Frenkel and N.Yu. Reshetikhin, “ Quantum affine algebras and holonomic difference equations.” Preprint (1991).
2. A. Tsuchiya and Y. Kanie, Advanced Studies in Pure Math. **16** (1988) 297, Lett. Math. Phys. **13** (1987) 303.
3. E. Date, M. Jimbo and M. Okado, “ Crystal base and q -vertex operators ” Osaka Univ. Math. Sci. preprint 1 (1991).
4. B. Davies, O. Foda, M. Jimbo, T. Miwa and A. Nakayashiki, “ Diagonalization of the XXZ hamiltonian by vertex operators ” RIMS preprint (1992).
5. H. Awata and Y. Yamada, “ Fusion rules for the fractional level $\widehat{sl}(2)$ algebra ” KEK Preprint 91-209 KEK-TH-316 (1992), to appear in Mod. Phys. Lett. **A**.
6. F.A. Smirnov, “ Dynamical symmetries of massive integrable models ” RIMS preprint 772, 838 (1991).
A. LeClair and F.A. Smirnov, “ Infinite quantum group symmetry of fields in massive 2D quantum field theory ” CLNS preprint 91-1056 (1991).
7. G. Felder and A. LeClair, “ Restricted quantum affine symmetry of perturbed minimal conformal models ” RIMS preprint 799 (1991).
8. M. Bauer and N. Sochen, “ Fusion and singular vectors in $A_1^{(1)}$ highest weight cyclic modules ” Saclay preprint (1992).
9. F.G. Malikov, “ Quantum groups: Singular vectors and BGG resolution ” RIMS Preprint 835 (1991).